

NASW-3950  
DRA/HQ

(NASA-CR-175491) COMETARY ATMOSPHERES:  
MODELING THE SPATIAL DISTRIBUTION OF  
OBSERVED NEUTRAL RADICALS Interim Report, 1  
Jun. 1984 - 31 Aug. 1984 (Atmospheric and  
Environmental Research) 4 p EC A02/HF A01

N85-21097

Unclas

G3/91 14393

Cometary Atmospheres:  
Modeling the Spatial Distribution of Observed Neutral Radicals

Michael R. Combi

Atmospheric and Environmental Research, Inc.  
840 Memorial Drive  
Cambridge, MA 02139

October 1984

Interim Report for the Period  
June 1, 1984 to August 31, 1984

Prepared for  
NASA Headquarters



TECHNICAL REPORT STANDARD TITLE PAGE

1. Report No.	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle  Cometary Atmospheres: Modeling the Spatial Distribution of Observed Neutral Radicals		5. Report Date October 1984	
		6. Performing Organization Code	
7. Author(s) Michael R. Combi		8. Performing Organization Report No.	
9. Performing Organization Name and Address  Atmospheric and Environmental Research, Inc. 840 Memorial Drive Cambridge, Massachusetts 02139		10. Work Unit No.	
		11. Contract or Grant No. NASW-3950	
12. Sponsoring Agency Name and Address  NASA Headquarters Headquarters Contract Division Washington, DC 20546		13. Type of Report and Period Covered Interim Report June - August 1984	
		14. Sponsoring Agency Code HW-2	
15. Supplementary Notes			
16. Abstract  The Monte Carlo particle-trajectory model for the spatial distribution of cometary radicals has been modified to include the heliocentric distance dependence of the parent molecule velocity, and the heliocentric velocity dependence for CN fluorescence and radiation pressure. The study of available data on the observed spatial distributions of cometary radicals has begun and a preliminary comparison of some newly published data with previous studies is discussed.			
17. Key Words (Selected by Author(s))  comets		18. Distribution Statement	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages	22. Price*

\*For sale by the Clearinghouse for Federal Scientific and Technical Information, Springfield, Virginia 22151.

## I. Program of Research for the First Quarter

Research activities during this first quarter have involved (1) several refinements and modifications to the computer code for the Monte Carlo particle trajectory model, and (2) preliminary comparison of recently published radical parent Haser scale length data with previously published data.

### 1. Model Updates

Whipple (1980) has shown that the outflow velocity of parent molecules is a slowly decreasing function of heliocentric distance. Delsemme (1982) has derived a semi-empirical law for this dependence to be  $V = .58 r_H^{-1/2}$ , where  $V$  is in km/s and  $r_H$  is in astronomical units. This velocity dependence is now automatically incorporated into the computer code for the Monte Carlo model. It has been common practice to assume an  $r_H^2$  law for the parent molecule scale lengths in Haser's model. Observational evidence, on the other hand, suggests that a value for the exponent somewhat less than 2 may be more appropriate (Combi and Delsemme 1980), as would be expected from the above velocity dependence combined with an  $r_H^2$  dependence from photochemical lifetimes.

Tatum and Gillespie (1977) calculated the dependence on heliocentric velocity of the resonance fluorescence emission rate for the CN(0-0) violet bands at 3883 Å. The velocity dependence results from the variable (with Doppler shift) overlap of rotational lines of cometary CN with absorption lines in the solar spectrum, some of which are from the same CN band. Their variable emission rate data has been directly incorporated in the model since it affects both the observed emission rate as well as the radiation pressure acceleration.

In addition to these changes to the physics of the model, a computer code shell has been constructed around the model so as to enable the direct calculation of radical production rates from observed photometric band emission fluxes.

### 2. New Haser Scale Lengths

Newburn and Spinrad (1984) have recently published the results of spectrophotometry of 17 comets. In particular, they presented Haser scale lengths for the parent molecules of  $C_2$ ,  $C_3$  and CN for 5 of these comets.

A preliminary comparison of their data for CN with previously published CN scale lengths (Combi and Delsemme 1980; Delsemme and Combi 1983) is quite interesting. Their data covers a range of heliocentric distances from .78 to 1.69 AU which is quite complementary to our data which covered from .44 to 1.21 AU. Considering the large amount of scatter in this type of analysis, the two data sets seem to merge reasonably well with no gross systematic differences. However, comparison of their C<sub>2</sub> parent scale lengths with both previously published as well as our as yet unpublished values does show some apparent systematic differences. Since both the types of analyses as well as the two populations of comets studied were quite different, a more careful comparison of these two data sets is needed.

## II. Program of Research for the Next Quarter

Research activities during the next quarter will concentrate on (1) incorporation of the OH heliocentric velocity dependent lifetime and emission rate in the model, (2) continued study of available data on the spatial distributions of cometary radicals, and (3) a preliminary evaluation of the effect of radiation pressure on photometrically determined production rates.

## References

Combi, M.R. and A.H. Delsemme (1980), Ap. J. 237, 641.

Delsemme, A.H. (1982), Comets, Edited by Laurel L. Wilkening, The University of Arizona Press, Tucson, AZ, p. 85.

Delsemme, A.H. and M.R. Combi (1983), Ap. J. 271, 388.

Newburn, R.L. and H. Spinrad (1984), A. J. 89, 289.

Whipple, F.L. (1980), A. J. 85, 305.